

Coherent states of non-relativistic electron in magnetic – solenoid field

V. G. Bagrov*, S. P. Gavrilov†, D. M. Gitman‡, D. P. Meira Filho§

July 8, 2010

Abstract

In the present work we construct coherent states in the magnetic-solenoid field, which is a superposition of the Aharonov-Bohm field and a collinear uniform magnetic field. In the problem under consideration there are two kind of coherent states, those which correspond classical trajectories which embrace the solenoid and those which do not. Constructed coherent states reproduce exactly classical trajectories, they maintain their form under the time evolution, and form a complete set of functions, which can be useful in semiclassical calculations. In the absence of the solenoid field these states are reduced to well known in the case of a uniform magnetic field Malkin-Man'ko coherent states.

PACS numbers: 03.65.Ge, 03.65.Sq

1 Introduction

It is well-known that the study of the Aharonov-Bohm (AB) effect began on the base of exact wave functions of an electron in the field of an infinitely long and infinitesimally thin solenoid [1]. Such functions allow one to analyze a nontrivial influence of the AB solenoid (ABS) on scattering of free electron, which may give a new interpretation of electromagnetic potentials in quantum theory. Physically it is clear, that in such a scattering, the electron is subjected to the action of the AB field for a short finite time. However, there exist a possibility to consider bound states of the electron in which it is affected by the AB field for the infinite time. Such bound states exist in the so-called magnetic-solenoid field (MSF), which is a superposition of the AB field and a collinear uniform magnetic field. The non-relativistic and relativistic wave functions of an electron in the MSF were studied in [2, 3, 4, 5, 6]. In particular, they were used to describe AB effect in cyclotron and synchrotron radiations [7]. We believe that such bound states of an electron in the MSF open new possibilities the study of the AB effect. One of the important questions is the construction and study of semiclassical (coherent) states in the MSF (the importance and advantage of coherent states in quantum theory is well-known [8]). Having such states in hands one can try to answer an important question: to what extent the AB effect is of a pure quantum nature. In a sense constructing semiclassical states is a complementary task to the path integral construction. The latter problem is completely open in the case of the particle in the magnetic-solenoid field. One ought to say that some attempts to construct semiclassical states in the MSF were made in the works [9]. However, one ought to accept that states constructed in these works

*Department of Physics, Tomsk State University, 634050, Tomsk, Russia. Tomsk Institute of High Current Electronics, SB RAS, 634034 Tomsk, Russia; e-mail: bagrov@phys.tsu.ru

†Institute of Physics, University of São Paulo, Brazil; On leave from Department of general and experimental physics, Herzen State Pedagogical University of Russia, Moyka emb. 48, 191186 St. Petersburg, Russia; e-mail: gavrilovsergeyp@yahoo.com

‡Institute of Physics, University of São Paulo, CP 66318, CEP 05315-970 São Paulo, SP, Brazil; e-mail: gitman@dfn.if.usp.br

§Institute of Physics, University of São Paulo, Brazil; e-mail: dmeira@dfn.if.usp.br

have some features that does not allow one to interpret them as semiclassical and coherent states. For example, some mean values calculated in such states do not move along classical trajectories. In the present work we succeeded to construct another kind of semiclassical states in the MSF which can be really treated as coherent states. The progress is related to a nontrivial observation that in the problem under consideration there are two kind of coherent states those which correspond classical trajectories which embrace the solenoid and those which do not. Constructed coherent states reproduce classical trajectories in the semiclassical limit, they maintain their form under the time evolution, and form a complete set of functions, which can be useful in semiclassical calculations. In the absence of the AB field these states are reduced to well known in the case of a uniform magnetic field Malkin-Man'ko coherent states [10].

We consider the non-relativistic motion of an electron with charge $q = -e$, $e > 0$, and mass M in the MSF $\mathbf{B} = (B_x, B_y, B_z)$,

$$B_x = B_y = 0, \quad B_z = B + \Phi \delta(x) \delta(y) = B + \frac{\Phi}{\pi r} \delta(r), \quad (1)$$

which is a collinear superposition of a constant uniform magnetic field B directed along the axis z ($B > 0$) and the AB field (field of an infinitely long and infinitesimally thin solenoid) with a finite constant internal magnetic flux Φ . We use Cartesian coordinates x, y, z , as well as cylindrical coordinates r, φ , such that $x = r \cos \varphi$, $y = r \sin \varphi$, and $r^2 = x^2 + y^2$. The field (1) can be described by the vector potential $\mathbf{A} = (A_x, A_y, A_z)$,

$$A_x = -y \left(\frac{\Phi}{2\pi r^2} + \frac{B}{2} \right), \quad A_y = x \left(\frac{\Phi}{2\pi r^2} + \frac{B}{2} \right), \quad A_z = 0. \quad (2)$$

Classical motion of the electron in the MSF is governed by the Hamiltonian $H = \mathbf{P}^2/2M$, $\mathbf{P} = \mathbf{p} - \frac{q}{c}\mathbf{A}$, where $\mathbf{p}$ and $\mathbf{P}$ are the generalized and kinetic momentum, respectively. Trajectories that do not intersect the solenoid have the form:

$$x = x_0 + R \cos \psi, \quad y = y_0 + R \sin \psi, \quad z = \frac{p_z}{M}t + z_0; \quad \psi = \omega t + \psi_0, \quad \omega = \frac{eB}{Mc}, \quad (3)$$

where x_0, y_0, z_0, p_z, R , and ψ_0 are integration constants. Eqs. (3) imply

$$(x - x_0)^2 + (y - y_0)^2 = R^2, \quad x_0 = R_c \cos \alpha, \quad y_0 = R_c \sin \alpha, \\ r^2 = R^2 + R_c^2 + 2RR_c \cos(\psi - \alpha), \quad R_c = \sqrt{x_0^2 + y_0^2}. \quad (4)$$

The projection of particle trajectories on the xy -plane are circles. Particle images on the xy -plane are rotating with the synchrotron frequency ω . For an observer which is placed near the solenoid with $z > 0$, the rotation is anticlockwise. The particle has a constant velocity p_z/M along the axis z . Since the electron freely propagates on the z axis, only motion in the perpendicular plane $z = 0$ is nontrivial; this will be examined below. Denoting by r_{max} the maximal possible moving off and by r_{min} the minimal possible moving off of the particle from the z -axis, we obtain from (4) $r_{max} = R + R_c$, $r_{min} = |R - R_c|$. It follows from (3) that

$$P_x = -M\omega R \sin \psi = -M\omega (y - y_0), \\ P_y = M\omega R \cos \psi = M\omega (x - x_0), \quad \mathbf{P}_\perp^2 = P_x^2 + P_y^2 = (M\omega R)^2. \quad (5)$$

The energy E of the particle rotation reads: $E = \mathbf{P}_\perp^2/2M$ then the radius R can be expressed via the energy E as follows

$$R^2 = \frac{2E}{M\omega^2}. \quad (6)$$




Figure 1: Two types of trajectories in MSF

By the help of (5), one can calculate angular momentum projection L_z ,

$$L_z = xp_y - yp_x = \frac{M\omega}{2}(R^2 - R_c^2) - \frac{e\Phi}{2\pi c} . \quad (7)$$

The presence of ABS breaks the translational symmetry in the xy -plane, which on the classical level has only a topological effect, there appear two types of trajectories, we label them by an index $j = 0, 1$ in what follows. On the classical level $j = 1$ corresponds to $(R^2 - R_c^2) > 0$ and $j = 0$ corresponds to $(R^2 - R_c^2) < 0$, see Fig. 1.

Already in classical theory, it is convenient, to introduce dimensionless complex quantities a_1 and a_2 (containing $\hbar$) as follows:

$$a_1 = \frac{-iP_x - P_y}{\sqrt{2\hbar M\omega}} = -\sqrt{\frac{M\omega}{2\hbar}} R e^{-i\psi}, \quad a_2 = \frac{M\omega(x + iy) + iP_x - P_y}{\sqrt{2\hbar M\omega}} = \sqrt{\frac{M\omega}{2\hbar}} R_c e^{i\alpha}. \quad (8)$$

One can see that $a_1 \exp(i\psi)$ and a_2 are complex (dependent) integrals of motion. One can write that

$$R^2 = \frac{2\hbar}{M\omega} a_1^* a_1, \quad R_c^2 = \frac{2\hbar}{M\omega} a_2^* a_2, \quad x + iy = \sqrt{\frac{2\hbar}{M\omega}} (a_2 - a_1^*), \quad (9)$$

$$E = \omega \hbar a_1^* a_1, \quad L_z = \hbar (a_1^* a_1 - a_2^* a_2) - \frac{e\Phi}{2\pi c} . \quad (10)$$

2 Stationary states

The quantum behavior of the electron in the field (1) is determined by the Schrödinger equation with the Hamiltonian

$$\hat{H} = \hat{H}_\perp + \hat{p}_z^2/2M, \quad \hat{H}_\perp = (\hat{P}_x^2 + \hat{P}_y^2)/2M, \\ \hat{P}_x = \hat{p}_x + \frac{e}{c} A_x, \quad \hat{P}_y = \hat{p}_y + \frac{e}{c} A_y, \quad \hat{p}_x = -i\hbar\partial_x, \quad \hat{p}_y = -i\hbar\partial_y, \quad \hat{p}_z = -i\hbar\partial_z, \quad (11)$$

where $\hat{H}_\perp$ determines the nontrivial behavior on the xy -plane. It is convenient to present magnetic flux Φ in eq. (2) as $\Phi = (l_0 + \mu)\Phi_0$, where l_0 is integer, and $0 \leq \mu < 1$ and

$\Phi_0 = 2\pi\hbar/e$ is Dirac's fundamental unit of magnetic flux. Mantissa of the magnetic flux μ determines, in fact, all the quantum effects due to the presence of the AB field. Stationary states of the non-relativistic electron in the MSF were first described in [2]. The corresponding radial functions were taken regular at $r \rightarrow 0$, they correspond to a most natural self-adjoint extension (with a domain $D_{H_\perp}$) of the differential symmetric operator $\hat{H}_\perp$. Considering a regularized case of a finite-radius solenoid one can demonstrate that the zero-radius limit yields such an extension, see [6]. Further, we consider only such an extension (all possible self-adjoint extensions of $\hat{H}_\perp$ were constructed in ([4, 5])). Operator $\hat{L}_z = x\hat{p}_y - y\hat{p}_x$ is self-adjoint on $D_{H_\perp}$ and commutes with the self-adjoint Hamiltonian $\hat{H}_\perp$. One can find two types ($j = 0, 1$) of common eigenfunctions of both operators

$$\begin{aligned}\hat{H}_\perp \Psi_{n_1, n_2}^{(j)}(t, r, \varphi) &= \mathcal{E}_{n_1} \Psi_{n_1, n_2}^{(j)}(t, r, \varphi), \quad \mathcal{E}_{n_1} = \hbar\omega(n_1 + 1/2), \\ \hat{L}_z \Psi_{n_1, n_2}^{(j)}(t, r, \varphi) &= L_z \Psi_{n_1, n_2}^{(j)}(t, r, \varphi), \quad L_z = \hbar(l - l_0).\end{aligned}\quad (12)$$

The eigenfunctions have the form

$$\begin{aligned}\Psi_{n_1, n_2}^{(j)}(t, r, \varphi) &= \exp\left(-\frac{i}{\hbar}\mathcal{E}_{n_1}t\right) \Phi_{n_1, n_2}^{(j)}(\varphi, \rho), \quad \rho = \frac{eBr^2}{2c\hbar}, \quad j = 0, 1, \\ \Phi_{n_1, n_2}^{(0)}(\varphi, \rho) &= \mathcal{N} \exp[i(l - l_0)\varphi] I_{n_2, n_1}(\rho), \quad n_1 = m, \quad n_2 = m - l - \mu, \quad -\infty < l \leq -1, \\ \Phi_{n_1, n_2}^{(1)}(\varphi, \rho) &= \mathcal{N} \exp[i(l - l_0)\varphi - i\pi l] I_{n_1, n_2}(\rho), \quad n_1 = m + l + \mu, \quad n_2 = m, \quad 0 \leq l \leq +\infty.\end{aligned}\quad (13)$$

Here l, m ($m \geq 0$) are two integers, $I_{n, m}(\rho)$ are Laguerre functions that are related to the Laguerre polynomials $L_m^\alpha(\rho)$ (see eqs. 8.970, 8.972.1 from [11]) as follows

$$I_{m+\alpha, m}(\rho) = \sqrt{\frac{\Gamma(m+1)}{\Gamma(m+\alpha+1)}} e^{-\rho/2} \rho^{\alpha/2} L_m^\alpha(\rho), \quad L_m^\alpha(\rho) = \frac{1}{m!} e^\rho \rho^{-\alpha} \frac{d^m}{d\rho^m} e^{-\rho} \rho^{m+\alpha}, \quad (14)$$

and $\mathcal{N}$ is normalization constant. For any real $\alpha > -1$ the functions $I_{\alpha+m, m}(\rho)$ form a complete orthonormal set on the half-line $\rho \geq 0$,

$$\int_0^\infty I_{\alpha+k, k}(\rho) I_{\alpha+m, m}(\rho) d\rho = \delta_{k, m}, \quad \sum_{m=0}^\infty I_{\alpha+m, m}(\rho) I_{\alpha+m, m}(\rho') = \delta(\rho - \rho'). \quad (15)$$

Let us define an inner product of two functions $f(\varphi, \rho)$ and $g(\varphi, \rho)$ as

$$(f, g)_\perp = \frac{1}{2\pi} \int_0^\infty d\rho \int_0^{2\pi} d\varphi f^*(\varphi, \rho) g(\varphi, \rho).$$

Then eigenfunctions (13) form an orthogonal set on the xy -plane,

$$\left(\Psi_{n'_1, n'_2}^{(j')}, \Psi_{n_1, n_2}^{(j)}\right)_\perp = |\mathcal{N}|^2 \delta_{n'_1, n_1} \delta_{n'_2, n_2} \delta_{j', j}. \quad (16)$$

These functions form a complete orthogonalized set on $D_{H_\perp}$.

It is useful to define self-adjoint operators $\hat{R}^2$ and $\hat{R}_c^2$ by analogy with the corresponding classical relations (6) and (7):

$$\hat{R}^2 = \frac{2\hat{H}_\perp}{M\omega^2}, \quad \hat{R}_c^2 = \hat{R}^2 - \frac{2}{M\omega} \left[\hat{L}_z + (l_0 + \mu)\hbar \right]. \quad (17)$$

In the semiclassical limit the sign of the mean value of the operator $\hat{R}^2 - \hat{R}_c^2$,

$$\left(\Psi_{n_1, n_2}^{(j)}, \left(\hat{R}^2 - \hat{R}_c^2\right) \Psi_{n_1, n_2}^{(j)}\right)_\perp |\mathcal{N}|^{-2} = \frac{2\hbar(l + \mu)}{M\omega},$$




Figure 2: Splitting of Landau levels in MSF.

where (12) is used, allows one to interpret the corresponding states as particle trajectories that embrace and do not embrace the solenoid. Namely, an orbit embraces the solenoid for $l \geq 0$ (type $j = 1$), and do not for $l \leq -1$ (type $j = 0$). These classification corresponds to classical one described in the previous section, see eq. (7) and Fig. 1. Trajectories with $l = 0$ and $l = -1$ pass most close to the solenoid.

If $\mu \neq 0$, the degeneracy of energy spectrum is partially lifted, namely, energy levels of states (13) with $l \geq 0$ are shifted with respect to the Landau levels by $\mu\hbar\omega$, such that $\mathcal{E}_{n_1} = \hbar\omega(m + l + \mu + 1/2)$, while energy levels of states (13) with $l \leq -1$ are still given by the Landau formula $\mathcal{E}_{n_1} = \hbar\omega(m + 1/2)$. For $\mu = 0$ there is no any impact of ABS on the energy spectrum. Splitting of the Landau levels in the MSF is represented on Fig. 2.

3 Coherent states

3.1 Instantaneous coherent states on xy -plane

Let us introduce operators $\hat{a}_1$, $\hat{a}_2$ and $\hat{a}_1^\dagger, \hat{a}_2^\dagger$ that correspond to classical quantities a_1 , a_2 and a_1^* , a_2^* ,

$$\begin{aligned} \hat{a}_1 &= \frac{-i\hat{P}_x - \hat{P}_y}{\sqrt{2\hbar M\omega}}, \quad \hat{a}_2 = \frac{M\omega(x + iy) + i\hat{P}_x - \hat{P}_y}{\sqrt{2\hbar M\omega}}, \\ \hat{a}_1^\dagger &= \frac{i\hat{P}_x - \hat{P}_y}{\sqrt{2\hbar M\omega}}, \quad \hat{a}_2^\dagger = \frac{M\omega(x - iy) - i\hat{P}_x - \hat{P}_y}{\sqrt{2\hbar M\omega}}. \end{aligned} \quad (18)$$

One ought to say that the momentum operators $\hat{P}_x$ and $\hat{P}_y$ are symmetric but not self-adjoint on the domain $D_{H_\perp}$. That is why, one cannot consider $\hat{a}_1^\dagger$ and $\hat{a}_2^\dagger$ as adjoint to $\hat{a}_1$ and $\hat{a}_2$. Nevertheless, they play an important auxiliary role in the further constructions.

Using properties of Laguerre functions, one can find the action of the operators $\hat{a}_1^\dagger$, $\hat{a}_1$; $\hat{a}_2^\dagger$, $\hat{a}_2$ on the functions (13),

$$\begin{aligned} \hat{a}_1 \Phi_{n_1, n_2}^{(j)}(\varphi, \rho) &= \sqrt{n_1} \Phi_{n_1-1, n_2}^{(j)}(\varphi, \rho), \quad \hat{a}_1^\dagger \Phi_{n_1, n_2}^{(j)}(\varphi, \rho) = \sqrt{n_1+1} \Phi_{n_1+1, n_2}^{(j)}(\varphi, \rho), \\ \hat{a}_2 \Phi_{n_1, n_2}^{(j)}(\varphi, \rho) &= \sqrt{n_2} \Phi_{n_1, n_2-1}^{(j)}(\varphi, \rho), \quad \hat{a}_2^\dagger \Phi_{n_1, n_2}^{(j)}(\varphi, \rho) = \sqrt{n_2+1} \Phi_{n_1, n_2+1}^{(j)}(\varphi, \rho), \end{aligned} \quad (19)$$

where possible values of n_1 and n_2 depend on m, l , and j according to (13) and the functions

$\Phi_{n_1+s_1, n_2+s_2}^{(j)}$ are defined as follows

$$\begin{aligned}\Phi_{n_1+s_1, n_2+s_2}^{(0)}(\varphi, \rho) &= \mathcal{N} \exp[i\epsilon(l_0 - l - s_1 + s_2)\varphi] I_{n_2+s_2, n_1+s_1}(\rho), \\ \Phi_{n_1+s_1, n_2+s_2}^{(1)}(\varphi, \rho) &= \mathcal{N} \exp\{i\epsilon[(l_0 - l - s_1 + s_2)\varphi + \pi(l + s_1 - s_2)]\} I_{n_1+s_1, n_2+s_2}(\rho).\end{aligned}$$

at $s_1 = 0, \pm 1$ and $s_2 = 0, \pm 1$. There appear new functions $\Phi_{n_1, n_2-1}^{(0)}(\varphi, \rho)$ with $n_2 = m+1-\mu$ and $\Phi_{n_1-1, n_2}^{(1)}(\varphi, \rho)$ with $n_1 = m+\mu$, which are irregular at $r \rightarrow 0$. Such functions were not defined by eqs. (13). In addition, for $n_1 = 0$ or $n_2 = 0$, one has to bear in mind that

$$\hat{a}_1 \Phi_{0, -l-\mu}^{(0)}(\varphi, \rho) = 0, \quad \hat{a}_2 \Phi_{l+\mu, 0}^{(1)}(\varphi, \rho) = 0.$$

Formal commutators for the operators $\hat{a}_1^\dagger, \hat{a}_1$ and $\hat{a}_2^\dagger, \hat{a}_2$ have the form:

$$[\hat{a}_1, \hat{a}_1^\dagger] = 1 + f, \quad [\hat{a}_2, \hat{a}_2^\dagger] = 1 - f, \quad [\hat{a}_1, \hat{a}_2] = f, \quad [\hat{a}_1, \hat{a}_2^\dagger] = 0, \quad (20)$$

with a singular function $f = \Phi(\pi Br)^{-1} \delta(r) = 2(l_0 + \mu)\delta(\rho)$. However, one can verify by the help of (19) that this function gives zero contribution on the domain $D_{H_\perp}$, such that on this domain $\hat{a}_1^\dagger, \hat{a}_2^\dagger$ and $\hat{a}_1, \hat{a}_2$ behave as creation and annihilation operators. Operators $\hat{R}^2, \hat{R}_c^2, \hat{r}^2, \hat{H}_\perp$, and $\hat{L}_z$ can be expressed in terms of the operators $\hat{a}_1^\dagger, \hat{a}_1$ and $\hat{a}_2^\dagger, \hat{a}_2$ as follows:

$$\begin{aligned}\hat{R}^2 &= \frac{\hbar}{M\omega} (2\hat{N}_1 + 1), \quad \hat{R}_c^2 = \frac{\hbar}{M\omega} (2\hat{N}_2 + 1), \quad x + iy = \sqrt{\frac{2\hbar}{M\omega}} (\hat{a}_2 - \hat{a}_1^\dagger), \\ \hat{H}_\perp &= \hbar\omega (\hat{N}_1 + 1/2), \quad \frac{1}{\hbar} \hat{L}_z + l_0 + \mu = (\hat{N}_1 - \hat{N}_2), \quad \hat{N}_s = \hat{a}_s^\dagger \hat{a}_s, \quad s = 1, 2.\end{aligned} \quad (21)$$

The functions $\Phi_{n_1, n_2}^{(j)}(\varphi, \rho)$ (13) can be used to construct the following useful states $\Phi_{z_1, z_2}^{(j)}(\varphi, \rho)$,

$$\Phi_{z_1, z_2}^{(j)}(\varphi, \rho) = \sum_l \Phi_{z_1, z_2}^{(j)l}(\varphi, \rho), \quad \Phi_{z_1, z_2}^{(j)l}(\varphi, \rho) = \sum_m \frac{z_1^{n_1} z_2^{n_2} \Phi_{n_1, n_2}^{(j)}(\varphi, \rho)}{\sqrt{\Gamma(1+n_1) \Gamma(1+n_2)}}, \quad (22)$$

where z_1 and z_2 are complex parameters, possible values of n_1 and n_2 depend on m, l , and j according to (13), and we set $\mathcal{N} = 1$. We call these states instantaneous coherent states on xy -plane. These states can be expressed via special functions $Y_\alpha(z_1, z_2; \rho)$,

$$Y_\alpha(z_1, z_2; \rho) = \sum_{m=0}^{\infty} \frac{z_1^m z_2^{m+\alpha} I_{m+\alpha, m}(\rho)}{\sqrt{\Gamma(1+m) \Gamma(1+m+\alpha)}}, \quad (23)$$

as follows:

$$\begin{aligned}\Phi_{z_1, z_2}^{(0)l}(\varphi, \rho) &= \exp[i\epsilon(l_0 - l)\varphi] Y_{-l-\mu}(z_1, z_2; \rho), \\ \Phi_{z_1, z_2}^{(1)l}(\varphi, \rho) &= \exp\{i\epsilon[(l_0 - l)\varphi + \pi l]\} Y_{l+\mu}(z_2, z_1; \rho).\end{aligned} \quad (24)$$

By the help of the well-known sum,

$$\sum_{m=0}^{\infty} \frac{z^m I_{\alpha+m, m}(x)}{\sqrt{\Gamma(1+m) \Gamma(1+\alpha+m)}} = z^{-\frac{\alpha}{2}} \exp\left(z - \frac{x}{2}\right) J_\alpha(2\sqrt{xz}),$$

where $J_\alpha(x)$ are the Bessel functions of the first kind, one can obtain the following representation for $Y_\alpha(z_1, z_2; \rho)$:

$$Y_\alpha(z_1, z_2; \rho) = \exp\left(z_1 z_2 - \frac{\rho}{2}\right) \left(\sqrt{\frac{z_2}{z_1}}\right)^\alpha J_\alpha(2\sqrt{z_1 z_2} \rho). \quad (25)$$

Then it follows from (19):

$$\hat{N}_k \Phi_{z_1, z_2}^{(j)}(\varphi, \rho) = z_k \partial_{z_k} \Phi_{z_1, z_2}^{(j)}(\varphi, \rho), \quad k = 1, 2; \quad (26)$$

and

$$\begin{aligned} a_1 \Phi_{z_1, z_2}^{(j)}(\varphi, \rho) &= z_1 \left[\Phi_{z_1, z_2}^{(j)}(\varphi, \rho) - (-1)^j \Phi_{z_1, z_2}^{(j)-1}(\varphi, \rho) \right], \\ a_2 \Phi_{z_1, z_2}^{(j)}(\varphi, \rho) &= z_2 \left[\Phi_{z_1, z_2}^{(j)}(\varphi, \rho) + (-1)^j \Phi_{z_1, z_2}^{(j)0}(\varphi, \rho) \right]. \end{aligned} \quad (27)$$

Then, using eqs. 6.615 from [11], we obtain:

$$\begin{aligned} \left(\Phi_{z_1, z_2}^{(j)}, \Phi_{z'_1, z'_2}^{(j')} \right) &= \delta_{jj'} \mathcal{R}^{(j)}; \\ \mathcal{R}^{(0)} &= Q_{1-\mu} \left(\sqrt{z_1^* z'_1}, \sqrt{z_2^* z'_2} \right), \quad \mathcal{R}^{(1)} = Q_\mu \left(\sqrt{z_2^* z'_2}, \sqrt{z_1^* z'_1} \right), \\ Q_\alpha(u, v) &= Q_\alpha^-(u, v) + \left(\frac{v}{u} \right)^\alpha I_\alpha(2uv), \quad Q_\alpha^-(u, v) = \sum_{l=1}^{\infty} \left(\frac{v}{u} \right)^{\alpha+l} I_{\alpha+l}(2uv), \end{aligned} \quad (28)$$

where $I_\alpha(u)$ are the modified Bessel functions of the first kind. We define the mean value of an operator $\hat{F}$ in the form

$$\overline{(F)}_{(j)} = \left(\Phi_{z_1, z_2}^{(j)}, \hat{F} \Phi_{z_1, z_2}^{(j)} \right) \left(\Phi_{z_1, z_2}^{(j)}, \Phi_{z_1, z_2}^{(j)} \right)^{-1}.$$

Using (26), one can calculate the mean values of $\hat{N}_s$:

$$\overline{(N_s)}_{(j)} = z_s \left. \partial_{z'_s} \ln \mathcal{R}^{(j)} \right|_{z'_s = z_s}, \quad s = 1, 2. \quad (29)$$

This allows one to connect means values of $\hat{R}^2$ and $\hat{R}_c^2$ with parameters z_1 and z_2 . We expect in the semiclassical limit that $\overline{(N_s)}_{(j)} \approx |z_s|^2$. At the same time length scales defined by the means $\overline{(R^2)}_{(j)}$, $\overline{(R_c^2)}_{(j)}$ have to be large enough which implies $|z_s|^2 \gg 1$. We expect that the sign of the difference $\overline{(R^2)}_{(j)} - \overline{(R_c^2)}_{(j)}$ is related to the trajectory type if the difference is sufficiently large, such that for states with $j = 0$ we have $|z_1|^2 < |z_2|^2$, and for states with $j = 1$, we have $|z_1|^2 > |z_2|^2$. We note that in both cases the corresponding functions $Q_\alpha(u, v)$ are calculated at $|v| > |u| \gg 1$.

There exist all the derivatives $\partial_v \left[(v/u)^{\alpha+l} I_{\alpha+l}(2uv) \right]$, the series $Q_\alpha^-(u, v)$ converges and the series of derivatives $\sum_{l=1}^{\infty} \partial_v \left[(v/u)^{\alpha+l} I_{\alpha+l}(2uv) \right]$ converges uniformly on the half-line, $0 < \text{Re } v < \infty$. Thus, one arrives to a differential equation with respect to $Q_\alpha^-(u, v)$,

$$\frac{dQ_\alpha^-(u, v)}{dv} = 2v \left[(v/u)^\alpha I_\alpha(2uv) + Q_\alpha^-(u, v) \right].$$

To evaluate asymptotics, we represent its solution as follows:

$$Q_\alpha^-(u, v) = e^{u^2+v^2} [1 - T(u, v)], \quad T(u, v) = 2e^{-u^2} \int_v^\infty e^{-\tilde{v}^2} \left(\frac{\tilde{v}}{u} \right)^\alpha I_\alpha(2u\tilde{v}) \tilde{v} d\tilde{v}, \quad (30)$$

where formula 6.631.4 [11] is used. Then

$$Q_\alpha(u, v) = e^{u^2+v^2} \tilde{Q}_\alpha(u, v), \quad \tilde{Q}_\alpha(u, v) = \left[1 - T(u, v) + e^{-u^2-v^2} (v/u)^\alpha I_\alpha(2uv) \right]. \quad (31)$$

Thus, the mean values (29) have the form:

$$\begin{aligned} \overline{(N_s)}_{(j)} &= |z_s|^2 + z_s \partial_{z'_s} \ln \tilde{\mathcal{R}}^{(j)} \Big|_{z'_s=z_s}, \quad s = 1, 2, \\ \tilde{\mathcal{R}}^{(0)} &= \tilde{Q}_{1-\mu} \left(\sqrt{z_1^* z'_1}, \sqrt{z_2^* z'_2} \right), \quad \tilde{\mathcal{R}}^{(1)} = \tilde{Q}_\mu \left(\sqrt{z_2^* z'_2}, \sqrt{z_1^* z'_1} \right). \end{aligned} \quad (32)$$

Using asymptotics of the function $I_\alpha(2uv)$, one can verify that if $|v| > |u| \gg 1$ then $|z_s|^2 \gg z_s \partial_{z'_s} \ln \tilde{\mathcal{R}}^{(j)} \Big|_{z'_s=z_s}$ in (32). For semiclassical states corresponding to orbits placed far enough from the solenoid, i.e., for $||z_1|^2 - |z_2|^2| \gg 1$, the contribution $z_s \partial_{z'_s} \ln \tilde{\mathcal{R}}^{(j)} \Big|_{z'_s=z_s}$ is small as $\exp\left(-||z_1|^2 - |z_2|^2|\right)$. Finally, we obtain:

$$|z_1|^2 \approx \frac{M\omega}{2\hbar} \overline{(R^2)}_{(j)}, \quad |z_2|^2 \approx \frac{M\omega}{2\hbar} \overline{(R_c^2)}_{(j)}, \quad |z_s|^2 \gg 1. \quad (33)$$

By the help of (27), one can find:

$$\begin{aligned} \overline{(a_1)}_{(0)} &= z_1 \Delta_{1-\mu}(|z_1|, |z_2|), \quad \overline{(a_2)}_{(0)} = z_2, \quad \overline{(a_1)}_{(1)} = z_1, \\ \overline{(a_2)}_{(1)} &= z_2 \Delta_\mu(|z_2|, |z_1|), \quad \Delta_\alpha(u, v) = \frac{Q_\alpha^-(u, v)}{Q_\alpha(u, v)}, \end{aligned} \quad (34)$$

such that these means match with eqs. (32) in the classical limit.

3.2 Time-dependent coherent states

Consider Schrödinger equation with the complete three-dimensional Hamiltonian $\hat{H}$ (11) and corresponding solutions $\Psi(t, \mathbf{r})$ with a given momentum p_z ,

$$\Psi(t, \mathbf{r}) = \mathcal{N} \exp \left\{ -\frac{i}{\hbar} \left[\left(\frac{p_z^2}{2M} + \frac{\hbar\omega}{2} \right) t - p_z z \right] \right\} \Phi(t, \varphi, \rho),$$

where $\mathcal{N}$ is normalization constant. The functions $\Phi(t, \varphi, \rho)$ obey the following equation

$$i\partial_t \Phi(t, \varphi, \rho) = \omega \hat{N}_1 \Phi(t, \varphi, \rho). \quad (35)$$

One can obey (35) setting $\Phi(t, \varphi, \rho) = \Phi_{z_1, z_2}^{(j)}(\varphi, \rho) \Big|_{z_1=z_1(t)}$, where $z_1(t)$ is a complex function of time t . Then

$$i\partial_t \Psi_{z_1, z_2}^{(j)} = i\dot{z}_1 \partial_{z_1} \Phi_{z_1, z_2}^{(j)}, \quad \dot{z}_1 = dz_1/dt. \quad (36)$$

Substituting (36) into (35), we find $i\dot{z}_1 = \omega z_1$, where (26) is used. It is convenient to write a solution for $z_1(t)$ as follows:

$$z_1(t) = -|z_1| \exp(-i\psi), \quad \psi = \omega t + \psi_0, \quad (37)$$

where $|z_1|$ is a given constant. Thus the functions

$$\Psi_{\text{CS}}^{(j)}(t, \mathbf{r}) = \mathcal{N} \exp \left\{ -\frac{i}{\hbar} \left[\left(\frac{p_z^2}{2M} + \frac{\hbar\omega}{2} \right) t - p_z z \right] \right\} \Phi_{z_1(t), z_2}^{(j)}(\varphi, \rho) \quad (38)$$

are solutions of the Schrödinger equation. At the same time they have special properties that allow us to treat them as coherent (and under certain conditions as semiclassical) states.

Let us consider mean values $\overline{(x)}_{(j)}$ and $\overline{(y)}_{(j)}$ of the coordinates with respect to the states $\Psi_{\text{CS}}^{(j)}$. To this end it is enough to find the mean value $\overline{(x + iy)}_{(j)}$. By the help of (21) we obtain:

$$\overline{(x + iy)}_{(j)} = \sqrt{\frac{2\hbar}{M\omega}} \left[\overline{(a_2)}_{(j)} - \overline{(a_1)}_{(j)}^* \right].$$

Taking into account eqs. (34) and (37), one can see that a point with coordinates $\overline{(x)}_{(j)}$ and $\overline{(y)}_{(j)}$ is moving along a circle on the xy -plane with the cyclotron frequency ω , i.e., its trajectory has the classical form. The same equations allows one to find a radius $\overline{(R)}_{(j)}$ of such a circle and the distance $\overline{(R_c)}_{(j)}$ between its center and the origin,

$$\begin{aligned} \overline{(R)}_{(0)} &= \sqrt{\frac{2\hbar}{M\omega}} |z_1| \Delta_{1-\mu}(|z_1|, |z_2|), \quad \overline{(R_c)}_{(0)} = \sqrt{\frac{2\hbar}{M\omega}} |z_2|; \\ \overline{(R)}_{(1)} &= \sqrt{\frac{2\hbar}{M\omega}} |z_1|, \quad \overline{(R_c)}_{(1)} = \sqrt{\frac{2\hbar}{M\omega}} |z_2| \Delta_{\mu}(|z_2|, |z_1|). \end{aligned}$$

However, in the general case, the quantities $\overline{(R)}_{(j)}$ and $\overline{(R_c)}_{(j)}$ do not coincide with the corresponding quantities

$$\sqrt{\overline{(R^2)}_{(j)}} = \sqrt{\frac{\hbar}{M\omega}} \sqrt{2\overline{(N_1)}_{(j)} + 1}, \quad \sqrt{\overline{(R_c^2)}_{(j)}} = \sqrt{\frac{\hbar}{M\omega}} \sqrt{2\overline{(N_2)}_{(j)} + 1},$$

which are expressed in terms of mean values of the operators $\hat{H}_{\perp}$ and $\hat{L}_z$ according to (21), see also (29).

It follows from eq. (34) that $\Delta_{1-\mu}(|z_1|, |z_2|) < 1$ and $\Delta_{\mu}(|z_2|, |z_1|) < 1$. This allows us to give the following interpretation for two types of states with $j = 0, 1$. States with $j = 1$ correspond to orbits that embrace the ABS (which corresponds to $|z_1|^2 \gtrsim |z_2|^2$ in the semiclassical limit). For such orbits $\overline{(R_c)}_{(1)} < R_c$, where the quantity $R_c = \sqrt{2\hbar/M\omega} |z_2|$ is interpreted by us as a distance between ABS and the orbit center as a consequence of eq. (33). At the same time, the mean radius of the orbit coincides with the classical radius $R = \sqrt{2\hbar/M\omega} |z_1|$. The interpretation of R as the classical radius follows from eq. (33). States with $j = 0$ correspond to orbits that do not embraces the ABS (which corresponds to $|z_1|^2 \lesssim |z_2|^2$ in the semiclassical limit). For such orbits $\overline{(R_c)}_{(0)} = R_c$ and $\overline{(R)}_{(0)} < R$.

By using formulas (31), (32), (34) one can calculate the variances for $\hat{R}^2$, $\hat{R}_c^2$, and $x + y$ with respect to the coherent states in the semiclassical limit. With this result one can see these variances are relatively small for the semiclassical orbits situated far enough from the solenoid, i.e., for $\left| |z_1|^2 - |z_2|^2 \right| \gg 1$. In this case the coherent states are highly concentrated around the classical orbits. In the most interesting case when a semiclassical orbit is situated near the solenoid, such that the condition $\left| |z_1|^2 - |z_2|^2 \right| \ll 1$ holds, the variance for $x + y$ increases significantly while the variances for $\hat{R}^2$, $\hat{R}_c^2$ remain relatively small. In this case $R \approx R_c$, however, one has $\overline{(R_c)}_{(1)} < R$ and $\overline{(R)}_{(0)} < R_c$, as of course it must be for such semiclassical orbits. Having in mind that the standard deviation of $x + y$, δR , is relatively large at $R \approx R_c$, such that $\delta R \gg \left| R - \overline{(R_c)}_{(1)} \right|, \left| \overline{(R)}_{(0)} - R_c \right|$, we illustrate the typical spread of particle position around two types of semiclassical orbits at $R \approx R_c$ on Fig. 3.

Thus, for $\mu \neq 0$, classical relations between parameters of particle trajectory in constant magnetic field, such that relations between circle parameters R (related to particle energy) and R_c (related to particle angular momentum) are affected in the presence of ABS. Such relations do not feel the presence of ABS for $\mu = 0$, and, even for $\mu \neq 0$, in the classical limit (in the leading approximation for sufficiently large radii) discussed above.




Figure 3: Spread of particle position around two types of semiclassical orbits at $R \approx R_c$, where $\bar{R}_c = (\bar{R}_c)_{(1)}$ and $\bar{R} = (\bar{R})_{(0)}$.

Thus, in contrast to the problem in the constant uniform magnetic field (and in contrast to any problem with quadratic Hamiltonian) in the magnetic-solenoid field, we meet a completely new situation. Here time-dependent coherent states can be constructed (which is completely nontrivial fact due to nonquadratic nature of the Hamiltonian in the magnetic-solenoid field), the respective mean values move along classical trajectories, however classical relations between physical quantities imply additional semiclassical restrictions. Not all coherent states correspond to a semiclassical approximation, which is natural for nonquadratic Hamiltonians.

Finally, we ought to mention that only linear combinations of the form

$$\Psi(c_0, c_1; t, \mathbf{r}) = c_0 \Psi_{\text{CS}}^{(0)}(t, \mathbf{r}) + c_1 \Psi_{\text{CS}}^{(1)}(t, \mathbf{r}),$$

with c_0 and c_1 -arbitrary and $c_0 c_1 \neq 0$ were considered earlier as coherent states in [9]. Mean values of the operators $\hat{a}_1$ and $\hat{a}_2$ in such mixed states do not coincide with the classical expressions (8).

Acknowledgement V. G. Bagrov: this work is partially supported by Russian Science and Innovations Federal Agency under contract No 02.740.11.0238 and Russia President grant SS-3400.2010.2; S.P. Gavrilov thanks FAPESP for support and Universidade de São Paulo for hospitality; D.M. Gitman acknowledges the permanent support of FAPESP and CNPq; D. P. Meira Filho thanks CNPq for a support.

References

- [1] Aharonov Y and Bohm D 1959 Phys. Rev. **115** 485
- [2] Lewis R R 1983 Phys. Rev. **A28** 1228
- [3] Bagrov V G, Gitman D M and Tlyachev V B 2001 J. Math. Phys. **42** 1933; Falomir H and Pisani P A G 2001 J. Phys. A: Math. Gen **34** 4143; Bagrov V G, Baldiotti M C, Gitman D M and Shirokov I V 2002 J. Math. Phys. **43** 2284; Gavrilov S P, Gitman D M and Smirnov A A 2003 Phys. Rev. A **67** 024103; 2003 Eur. Phys. J. C **30** 009

- [4] Exner P, Št'oviček P and Vytrás P 2002 J. Math. Phys. **43** 2151
- [5] Gitman D M, Smirnov A, Tyutin I V, Voronov B L, arXiv:0911.0946 [quant-ph].
- [6] Gavrilov S P, Gitman D M, Smirnov A A and Voronov B L 2004 in: *Focus on Mathematical Physics Research* Ed. by Charles V. Benton (Nova Science, New York) p 131; hep-th/0308093.
- [7] Bagrov V G, Gitman D M, Levin A and Tlyachev V B 2001 Nucl. Phys. **B605** 425; 2001 Mod. Phys. Lett. **A16** 1171
- [8] Klauder J R, Sudarshan E C G 1968 *Fundamentals of Quantum Optics* (Benjamin, New York); Malkin I A, Man'ko V I 1979 *Dynamical Symmetries and Coherent States of Quantum Systems* (Nauka, Moscow); Klauder I R, Skagerstam B S 1985 *Coherent States, Applications in Physics and Mathematical Physics* (World Scientific, Singapore); Perelomov A M 1986 *Generalized Coherent States and Their Applications* (Springer, Berlin); Gazeau J-P 2009 *Coherent States in Quantum Optics* (Wiley-VCH, Berlin)
- [9] Bagrov V G, Gitman D M and Skarzhinsky V D 1986 Proc. Lebedev Inst. (Moscow) **176** 151; Bagrov V G, Gavrilov S P, Gitman D M and Meira Filho D P 2008 in: *Problems of Modern Theoretical Physics* Ed. V. Epp (Tomsk State University Press, Tomsk) p 57
- [10] Malkin I A and Man'ko V I 1968 Zh. Eksp. Teor. Fiz. **55** 1014
- [11] Gradshteyn I S and Ryzhik I M 2007 *Tables of Integrals, Series, and Products* (Academic Press, New York)